

Useful Telephone Numbers

- ENT Department, Lister Hospital
Telephone: 01438 314333, Ext. 5113 / 4118
- ENT Admission Office (waiting list)
Telephone: 01438 286836 or 01438 286835
- ENT Nurse Specialist (Monday to Friday, 8.30am - 6pm)
Telephone: 01438 314333, bleep 1028
Mobile: 0778 534 3359 or 0787 639 0290

Useful ENT website: www.entuk.org

In an Emergency:

Dial 999 and request an ambulance or attend the nearest A&E of the following hospitals unless away from home

- **Lister Hospital A&E (SG1 4AB) - 01438 784732**
- **New QEII Urgent Care Centre, Welwyn Garden City - 01707 247549**
- **Luton and Dunstable**
- **Princess Alexandra Hospital, Harlow**

**Lister Hospital
Coreys Mill Lane
Stevenage
Hertfordshire
SG1 4AB
Tel: 01438 314333**

www.enherts-tr.nhs.uk


Date of publication: April 2014
Author: Liza Delfin
Reference: ENT07 Version: 05 (Mar 2018)
Review Date: November 2019
© East and North Hertfordshire NHS Trust

You can request this information in a different format or another language.

Patient Information

Tonsillectomy (Adults)

Ear, Nose and Throat Department



Tonsillectomy in adults

Surgical removal of the tonsils is known as tonsillectomy.

The tonsils are lymphoid tissue, similar to the lymph nodes (or glands) found in the neck, groin and armpits. Tonsils are the two masses at the back of the throat. The tonsils play a part in the development of immunity to infection. When a patient has tonsillitis, they complain of severe sore throat and have a temperature and swollen glands in the neck. The patient is usually too unwell to go to work.

Tonsillitis is common and is often treated with antibiotics. Sometimes the tonsils become persistently infected and patients suffer from frequent attacks of tonsillitis over a period of time (recurrent tonsillitis). The tonsils become very large so may affect the breathing and may cause snoring.

What are the benefits?

Prevent recurrent tonsillitis / sore throat.

What are the alternatives, including no treatment?

You will not always need to have your tonsils out. You may want to wait and see if the tonsil problem gets better by itself.

The doctor or nurse specialist should explain to you why he/she feels that the surgery is the best treatment. You may change your mind about the operation at any time. If you would like to have a second opinion about the treatment, you can ask your specialist who can arrange this or you can get in touch with your GP.

How is the procedure performed?

The operation to remove the tonsils is carried out by dissecting the tonsil from the muscle bed of the wall of the throat. This may be done with electric or steel instruments. Bleeding is controlled with silk thread, ties or with coagulation using heat.

Length of operation

The operation takes between 20-40 minutes.

Anaesthetic

During general anaesthesia, the anaesthetist will, with the aid of the anaesthetic drug, keep you in a state of unconsciousness.

Risks and complications

Risks of anaesthesia largely depend on the overall health, the nature of the operation and its seriousness. Minor complications can include some soreness or bruising at the site of the injection

in the arm or hand. Some patients experience a sore throat if a tube has been inserted into the windpipe to assist breathing. On rare occasions this may damage teeth. Some of the drugs used can cause muscle pains and some patients may feel sick or nauseous.

General complications of surgery

Bleeding is normally controlled at the time of surgery. Occasionally bleeding can occur within the first 24 hours and may require further surgery. There is a risk of bleeding up to 7-14 days after the operation. If this happens it is because of an infection which will usually respond to antibiotics but if the bleeding persists further surgery may be needed.

Infection occurs in 3% of cases. Raised temperature in the first 24 hours is rarely due to infection. A persistent raised temperature may be due to an infection at the site of surgery or due to chest or urinary infections.

Pain - Painkillers should be taken regularly to lessen the discomfort.

Blood transfusion

Blood transfusions are rarely necessary unless severe bleeding occurs after the operation.

What happens after the procedure?

The operation is commonly performed as a day case or one night stay in hospital. Either way, we will only let you go home when you are eating and drinking and feel well enough.

It is very important to resume normal diet starting the day after the operation, drink plenty of water. Chewing and swallowing increases salivary flows, reduces spasm of pharyngeal muscles and removes debris from tonsil beds; this will also relieve discomfort and reduce the risk of infection. Avoid smoking and alcohol for two weeks.

Time off work

10 -14 days off work.

Post-operative care and advice

Remain indoors during the first week; avoid contact with people with infections; avoid exposure to dust and smoky places.

If there is profuse bleeding, report immediately to the nearest A&E Department.