


# Patient Information

## Pelvic Floor Weakness

Urogynaecology




## Introduction

The purpose of this leaflet is to explain what the pelvic floor is, what it does and how to exercise your pelvic floor to strengthen it.

## What is the pelvic floor?

The pelvic floor is composed of a thin sheet of muscle fibres and associated connective tissue which spans the area underneath the pelvis between the pubic bone at the front and the sacrum or tail bone at the back.

See the diagram here:



## What does the pelvic floor do?

- The pelvic floor muscles contract (tighten) when you cough, sneeze or strain, helping to prevent the involuntary leakage of urine.
- They help to support the organs in your abdomen, especially when you are standing.
- They help to protect the pelvic organs from external damage.
- They help to hold the pelvic organs, such as the bladder, in the correct position.
- They help in the control of passing urine, gas, and bowel motions.
- They play a role in sexual function during intercourse.

## What causes weakness to the pelvic floor?

- Under use. Like all muscles, the pelvic floor muscles need exercise to work well. Regular and adequate intensive exercise of the pelvic floor muscles should be maintained all through life.
- Damage to the muscles during pregnancy and childbirth.
- The change in hormones associated with menopause might play a role in weakening the muscles but is still debatable.
- Decreased muscle tone with ageing.
- Damage to the muscles through long-term straining when constipated or associated with a chronic cough or obesity.

### **A weakness of the pelvic floor muscles may cause or worsen a number of problems, including:**

- **Stress urinary incontinence** - The complaint of involuntary loss of urine on effort or physical exertion, including sporting activities etc., or on sneezing or coughing.
- **Urge urinary incontinence** - The complaint of involuntary loss of urine associated with a sudden urgent need to pass urine.
- **Mixed urgency and stress incontinence** - The complaint of involuntary loss of urine associated with urgency and also with effort or physical exertion, or on sneezing or coughing.
- **Pelvic organ prolapse** - Leading to a bulging of the bladder or bowel, or uterus (womb) into the vagina or out of the vaginal entrance if more severe.
- A loss in sexual sensation or feeling of vaginal looseness.

## How to exercise your pelvic floor muscles

It is important to learn to do the exercises in the right way, and to check from time to time that you are still doing them correctly. Your health professional can help you to understand this by using diagrams, drawings and models.

- 1) In the upright position, sit comfortably with your feet and knees wide apart. If preferred, you can also lie down flat (supine position) with your legs slightly apart. Remember to keep breathing throughout and keep your stomach, leg and buttock muscles relaxed.
- 2) Imagine that you are trying to stop yourself from passing gas from the bowel and at the same time trying to stop the flow of urine from the bladder. You should feel a lifting and tightening around the vagina and anus.
- 3) Observing the muscles can be an effective way of checking they are working correctly. Lie propped on the bed; place a small mirror between your legs with your knees bent and apart. As you pull up the pelvic floor you should see your anus and entrance to the vagina lifting and drawing in.

### **Note:**

- If you notice the entrance to the vagina widening, you are pushing down and not pulling up, which is the opposite to tightening the pelvic floor muscle.
- If you leak urine or pass gas when doing the exercises, it is likely you are pushing down not pulling up.

Don't worry if you are unable to contract and relax adequately at first. Most patients will accomplish this if they are given some time by themselves at home to practice. It can take time to perfect!

## Exercise 1

- 1) Tighten your pelvic floor muscles maximally without using your buttocks or thigh muscles as described on page 4, point (2). Hold tight for as many seconds as you can (up to a maximum of 10 seconds).
- 2) Release the contraction and rest for at least the same time of the contraction. Repeat the 'tighten, hold and release' as many times as you can (up to a maximum of 8-12 times)

**This program builds up the endurance of the muscle group. In other words, the muscle will be able to work harder for longer.**

## Exercise 2

- 1) Now perform the pelvic floor exercise, but squeeze quickly and then let go. Rest 1-2 seconds before the next one. These are called quick contractions and will help your muscles react quickly when you laugh, cough, exercise or lift.
- 2) Aim to increase this number to 8-12 contractions, 3 times a day for at least 6 months. That's the easy bit!

The most difficult part of the program is remembering to do the exercises!

## Can anything else help?

If you are unable to contract your muscles, your health professional may suggest some additional treatments.

**Biofeedback** - This aims to help women identify and contract and relax selectively the pelvic floor muscles using signals from their own bodies. Methods involve using small sticky electrodes placed on the abdomen and buttock area or an intra-vaginal or intrarectal probe connected to a computer. The strength and length of the contraction you perform can then be seen on the computer screen.

**Electrical stimulation therapy** - Sometimes the pelvic floor muscles cannot (yet) contract due to nerve damage. Electrical stimulation will provide an artificial contraction of the pelvic floor muscles, so helping to strengthen them. This is not suitable for women who can already contract their pelvic floor muscles, even if those contractions are weak.

**Avoid constipation** – Constipation and straining will increase the ‘downward’ pressure of tissue within the vagina and therefore may create or worsen a prolapse. It may also cause a prolapse to reoccur once corrected.

**Correct a persistent cough** – Prolonged and persistent coughing should be investigated to identify a cause as it will apply constant pressure to the pelvic floor muscles.

## Summary

- Weak pelvic muscles often cause bladder control problems.
- Daily intensive exercises can strengthen pelvic muscles and improve bladder control.
- If you are unsure, ask your health professional whether you are squeezing the right muscles.
- Always tighten and contract your pelvic muscles before coughing, sneezing, jumping or lifting. This can help prevent involuntary loss of urine, gas or stool, or bulging down of your pelvic organs.
- Try to maintain your weight within healthy limits.

### **When will I see results from doing pelvic floor exercises?**

You may not feel your bladder control improve until after 3-6 weeks, but it may take up to 6 months to have improvement.

### **Don't give up!!**

It is important to maintain the strength of the muscles once you have completed an exercise programme.

Practicing your exercises twice a week, 3 sets of 8-12 contractions, 3 times a day should be sufficient. Try to incorporate these exercises into daily life activities. If you are not sure that you are doing the exercises correctly, ask your family doctor, nurse, physiotherapist or continence nurse to check for you.

## Contact details

Urogynaecology Department  
The Woodlands Unit (Pink zone)  
Lister Hospital  
Stevenage  
Hertfordshire SG1 4AB

Telephone: 01438 286172 or 01438 288478  
Monday to Friday, 8am - 4pm

## Further reading

International Urogynecological Association (IUGA)  
[www.iuga.org/?page=patientleaflets](http://www.iuga.org/?page=patientleaflets)

Date of publication: January 2020  
Author: A.Digweed  
Reference: Version: 01  
Review Date: January 2023  
© East and North Hertfordshire NHS Trust

[www.enherts-tr.nhs.uk](http://www.enherts-tr.nhs.uk)

You can request this  
information in a different  
format or another language.