

Patient Information

Allergic Reaction

Children's Services

Introduction

The purpose of this leaflet is to explain the different symptoms and treatment of allergic reaction.

Causes of an allergic reaction - MINOR

Allergic reactions can occur in response to a substance that the body recognizes as foreign. It can be caused by food, drugs, substances or chemicals, animal furs, plants and pollens, house dust mite or by anything.

Symptoms of an allergic reaction

- Skin urticarial rash (hives), dry and itchy/burning skin
- Swelling of eyes/face/lips (angiodema)
- Eyes red watery eyes, itchy eyes, puffiness around the eyes
- Nasal sneezing, runny nose or persistent nasal discharges, rhinitis
- Stomach stomach pain, vomiting, diarrhoea

Treatment of an allergic reaction

- Give antihistamine as prescribed by your doctor
- Observe your child for signs of improvement or deterioration

Symptoms of an anaphylactic reaction - MAJOR

- **Airway** difficulty breathing, coughing, wheezy, choking, noisy breathing and voice change
- **Total body** change of skin colour, floppiness, sense of impending doom/feeling scared, loss of consciousness

Treatment of anaphylactic reaction

- If your child has an EPIPEN , give immediately
- Call 999
- If asthmatic, give 10 puffs of Salbutamol
- If there is no improvement in 5-10 minutes, a second EPIPEN should be given, if available

Further information	
----------------------------	--

NHS 111 or Contact your GP **2** 111

NHS website - www.nhs.uk

Useful Telephone Numbers

Bluebell Ward, Lister Hospital Stevenage	🛱 01438 284008
Children's Emergency Department Lister Hospital	2 01438 284333
Urgent Care Centre, New QEII Welwyn Garden City	2 01707 247549
Community Children's Nursing Team	2 01438 288370

Date of publication: June 2007 Author: Paediatric Team Version: 06 (June 2020) Review Date: June 2023 © East and North Hertfordshire NHS Trust

www.enherts-tr.nhs.uk

You can request this information in a different format or another language.